

## Born Too Soon and Too Small in Nebraska

### In An Average Week in Nebraska<sup>(1)</sup>

**53**  
babies are  
born preterm

**9**  
babies are born  
very preterm


**32**  
babies are born  
low birthweight

**6**  
babies are born very  
low birthweight


### Preterm Birth Rates

2010-2013 Average

Percent of live births (93 Counties)



Value ranges are based on an approximately equal number of counties in each range.



### Preterm and Low Birthweight Births in Nebraska

	2003	2013	2020 US Objective
<b>Preterm<sup>(1)</sup></b>	<b>12.3%</b>	<b>10.6%</b>	<b>11.4%</b>
<b>Low Birthweight<sup>(1)</sup></b>	<b>6.9%</b>	<b>6.4%</b>	<b>7.8%</b>

- In 2013, there were 2,756 preterm births in Nebraska, representing 10.6% of live births.<sup>(1)</sup>
- Between 2003 and 2013, the rate of infants born preterm in Nebraska declined nearly 14%.<sup>(1)</sup>
- During 2011-2013 (average) in Nebraska, preterm birth rates were highest for women ages 40 and older (17.7%), followed by women under age 20 (11.7%), ages 30-39 (11.3%) and ages 20-29 (10.0%).<sup>(1)</sup>
- During 2011-2013 (average) in Nebraska, preterm birth rates were highest for black infants (15.6%), followed by Hispanics (12.6%), Native Americans (11.7%), Asians (10.5%) and whites (9.9%).<sup>(1)</sup>
- In 2005, the annual societal economic cost (medical, educational, and lost productivity) associated with preterm birth in the United States was at least \$26.2 billion.<sup>(2)</sup>

## Born Too Soon and Too Small in Nebraska

**March of Dimes National Prematurity Campaign:** The Mission of the March of Dimes is to improve the health of babies by preventing birth defects, premature birth, and infant mortality. Prematurity is the leading cause of newborn death and a major determinant of illness and disability among infants, including developmental delays, chronic respiratory problems, and vision and hearing impairment. To address this growing problem, the March of Dimes has committed to a multi-year national prematurity research, awareness, and education campaign. For more information contact the pregnancy & newborn health education center at [askus@marchofdimes.org](mailto:askus@marchofdimes.org) or visit [marchofdimes.org/mission/prematurity-campaign.aspx](http://marchofdimes.org/mission/prematurity-campaign.aspx).

### Risk Factors for Preterm Birth

While the specific causes of spontaneous preterm labor and delivery are largely unknown, research indicates that they are likely due to a complex interplay of multiple risk factors, as opposed to any single isolated risk factor. Approximately one quarter of all preterm births are the result of medical intervention, and the remainder occur spontaneously. <sup>(3)</sup>

- The most consistently identified risk factors include a history of preterm birth, current multifetal pregnancy, and some uterine and/or cervical abnormalities. <sup>(4)</sup>
- Other possible risk factors include: infection (especially genito-urinary), diabetes mellitus, hypertension, late or no prenatal care, smoking, alcohol and illicit drug use. <sup>(4)</sup>
- In 2013, about 1 in 13 babies (7.5% of live births) was late preterm in Nebraska. The rise in late preterm births has been linked to rising rates of early induction of labor and c-sections. <sup>(5)</sup>
- In Nebraska in 2013, 8.9% of singleton births were preterm, compared to 56.4% of multiple births. Multiple births represent 3.5% of live births in Nebraska. <sup>(1)</sup>
- Smoking is an important determinant of health and a significant factor contributing to preterm and low birthweight births. In 2013, 20.8% of women of childbearing age reported smoking in Nebraska. <sup>(6)</sup>
- While being obese does not directly cause preterm birth, it does increase rates of medical complications (e.g., hypertension, diabetes) that do contribute to preterm birth. In Nebraska, 25.2% of women of childbearing age were obese in 2013. <sup>(6)</sup>

#### Footnotes

- Preterm is less than 37 completed weeks of pregnancy. Late preterm is between 34 and 36 weeks gestation. Very preterm is less than 32 completed weeks.
- Low birthweight is less than 2500 grams (5 1/2 pounds). Very low birthweight is less than 1500 grams (3 1/3 pounds).
- 2020 rates based on U.S. Healthy People Objectives.
- All race categories exclude Hispanics.
- Women of childbearing age are defined as 18-44 years.
- Obesity is defined as a Body Mass Index of 30 or more.

#### Source

- 1 National Center for Health Statistics, final natality data.
- 2 Institute of Medicine. 2007. Preterm Birth: Causes, Consequences, and Prevention. National Academy Press, Washington, D.C.
- 3 Iams JD. The epidemiology of preterm birth. Clin Perinatol. 2003;30:651-54.
- 4 Prematurity risk factors compiled by March of Dimes available at [www.marchofdimes.org/pregnancy/preterm-labor-and-birth.aspx](http://www.marchofdimes.org/pregnancy/preterm-labor-and-birth.aspx).
- 5 Bettgowda VR, Dias T, Davidoff MJ, et al. The relationship between cesarean delivery and gestational age among US singleton births. Clin Perinatol. 2008;35:309-323.
- 6 Behavioral Risk Factor Surveillance System, Centers for Disease Control and Prevention.

### Local Chapter

#### Nebraska

11640 Arbor Street, Suite 102 ,

Omaha, NE 68144

Phone: (402) 496-7111

E-mail: [NE408@marchofdimes.org](mailto:NE408@marchofdimes.org)

Website: <http://www.marchofdimes.com/nebraska>